arXiv:1002.0619v4 [math.NT] 28 Aug 2010

RECIPROCITY LAWS FOR LEGENDRE SYMBOLS OF THE TYPE $\left(\frac{a+b\sqrt{m}}{p}\right)$ - LONG VERSION[1]

CONSTANTIN N. BELI

Note This paper is an announcement. We do not prove the main result, Theorem 2.5 (as well as Theorem 2.10). We only show how Theorem 2.5 can be used to produce reciprocity laws for Legendre symbols of the type $\left(\frac{a+b\sqrt{m}}{p}\right)$. Note that in the applications we won't use Theorem 2.10, which is stronger than Theorem 2.5. Therefore readers who are not interested in details may skip Theorem 2.10 and everything related to it. (Lemma 1.1 and the part of §2 following Theorem 2.5.)

1. Introduction and notations

Given $p>2$ a prime and $a,b,m\in\mathbb{Z}/p\mathbb{Z}$ such that $\left(\frac{m}{p}\right)=\left(\frac{a^2-mb^2}{p}\right)=1$, the Legendre's symbol $\left(\frac{a+b\sqrt{m}}{p}\right)$ is defined as $\left(\frac{a+b\alpha}{p}\right)$, where $\alpha\in\mathbb{Z}/p\mathbb{Z}$ satisfies $\alpha^2=m$. The definition is independent of the choice of the square root α of m because $(a+b\alpha)(a-b\alpha)=a^2-b^2m$ so $\left(\frac{a+b\alpha}{p}\right)\left(\frac{a-b\alpha}{p}\right)=\left(\frac{a^2-mb^2}{p}\right)=1$ so $\left(\frac{a+b\alpha}{p}\right)=\left(\frac{a-b\alpha}{p}\right)$. One may aslo define $\left(\frac{a+b\sqrt{m}}{p}\right)$ if $a^2-mb^2=0$ and $a\neq 0$. In this case the two radicals of m are $\alpha=\pm\frac{a}{b}$. If we take $\alpha=-\frac{a}{b}$ then $\left(\frac{a+b\alpha}{p}\right)=\left(\frac{0}{p}\right)$, which is not convenient. Hence we will take $\alpha=\frac{a}{b}$ and we get $\left(\frac{a+b\sqrt{m}}{p}\right)=\left(\frac{a+b\alpha}{p}\right)=\left(\frac{2a}{p}\right)$.

In the particular case when $a=0$, $b=1$ we get $\left(\frac{\sqrt{m}}{p}\right)$, defined when $\left(\frac{m}{p}\right)=\left(\frac{0^2-m\cdot 1^2}{p}\right)=1$, i.e. when $p\equiv 1\pmod 4$ and $\left(\frac{m}{p}\right)=1$. This coincides with the rational quartic residue symbol $\left(\frac{m}{p}\right)_4$.

---

[1] THIS PAPER CONTAINS SOME RATHER LENGHTY DISCUTIONS INVOLVING 2-ADIC HILBERT SYMBOLS. READERS WHO WANT TO SKIP THIS TECHNICAL PART MAY CONSIDER THE SHORT VERSION.

The Legendre symbol $\left(\frac{a+b\sqrt{m}}{p}\right)$ can also be defined when a, b, m are p-adic integers satisfying $\left(\frac{m}{p}\right) = \left(\frac{a^2-mb^2}{p}\right) = 1$ or $p \mid a^2 - mb^2$, $p \nmid a$.

For any prime p of $\mathbb{Q}$, including the archimedian prime $p = \infty$, we denote by $\left(\frac{\cdot,\cdot}{p}\right) : \mathbb{Q}_p^\times/(\mathbb{Q}_p^\times)^2 \times \mathbb{Q}_p^\times/(\mathbb{Q}_p^\times)^2 \to \{\pm 1\}$ the Hilbert symbol. (At $p = \infty$ we have $\mathbb{Q}_\infty = \mathbb{R}$.)

Similarly as for $\left(\frac{a+b\sqrt{m}}{p}\right)$ we can introduce the Hilbert symbol $\left(\frac{a+b\sqrt{m},c}{p}\right)$. It is defined for $a, b, c, m \in \mathbb{Q}_p$ satisfying $m \in (\mathbb{Q}_p^\times)^2$ and $\left(\frac{a^2-mb^2,c}{p}\right) = 1$. If $\pm\alpha \in \mathbb{Q}_p^\times$ are the two square roots of m then $\left(\frac{(a+b\alpha)(a-b\alpha),c}{p}\right) = \left(\frac{a^2-mb^2,c}{p}\right) = 1$ so $\left(\frac{a+b\alpha,c}{p}\right) = \left(\frac{a-b\alpha,c}{p}\right)$ so there is no ambiguity in this definition. If $a^2 - mb^2 = 0$ and $a \neq 0$, same as for $\left(\frac{a+b\sqrt{m}}{p}\right)$, we define $\left(\frac{a+b\sqrt{m},c}{p}\right) = \left(\frac{2a,c}{p}\right)$. Throughout this paper we will consider $\left(\frac{a+b\sqrt{m},c}{p}\right)$ only for $p = 2$.

Various mathematicians have obtained results involving Legendre symbols of the type $\left(\frac{a+b\sqrt{m}}{p}\right)$. Most of this results involve the biquadratic symbols $\left(\frac{m}{p}\right)_4$ and they are called biquadratic or quartic reciprocity laws. In this paper we will give some very general reciprocity laws, which generalize all the existing results. In §3 we show how many of these results can be obtained as a consequence of Theorem 2.5, which we only state in this paper.

Definition 1. *For any $A \in \mathbb{Q}^\times/(\mathbb{Q}^\times)^2$ we denote by $\overline{A}$ the only square-free integer such that $A = \overline{A}(\mathbb{Q}^\times)^2$,*

The algebras $T(V)$, $S(V)$ and $S'(V)$.

For convenience we denote by $V := \mathbb{Q}^\times/(\mathbb{Q}^\times)^2$ and for any prime p we denote $V_p := \mathbb{Q}_p^\times/(\mathbb{Q}_p^\times)^2$. In particular, $V_\infty = \mathbb{R}^\times/(\mathbb{R}^\times)^2$.

Now V and V_p are $\mathbb{Z}/2\mathbb{Z}$-vector spaces. Therefore we may consider the corresponding tensor algebras $T(V)$ and $T(V_p)$ and the symmetric algebras $S(V)$ and $S(V_p)$. Note that, while V has a multiplicative notation, $T(V)$ and $S(V)$ have an additive one. In order to prevent confusion for any $A_1, \ldots, A_n \in V$ we will denote by $A_1 \bullet A_2 \bullet \cdots \bullet A_n$ their product in $S^n(V)$ (instead of simply $A_1 \cdots A_n$). For example, the distributivity law on $S^2(V)$ will be written as $AB \bullet C = A \bullet C + B \bullet C$.

We also define the algebra

$$S'(V) = T(V)/\langle A_1 \otimes \cdots \otimes A_n - A_{\pi(1)} \otimes \cdots \otimes A_{\pi(n)} \mid \pi \in A_n \rangle.$$

(Same definition as for $S(V)$ but this time the products are invariant only under even permutations of the factors.)

We denote by $A_1\odot\cdots\odot A_n$ the image in $S'(V)$ of $A_1\otimes\cdots\otimes A_n\in T(V)$. We will only be concerned with the homogenous component of degree 3, $S'^3(V)=T^3(V)/\langle A\otimes B\otimes C-B\otimes C\otimes A\rangle$. We have $A\odot B\odot C=B\odot C\odot A=C\odot A\odot B$ and $B\odot A\odot C=A\odot C\odot B=C\odot B\odot A$. However $A\odot B\odot C\neq B\odot A\odot C$ (unless A,B,C are linearly dependent). Similarly we define $S'(V_p)$.

For any $\xi\in V$, $T(V)$, $S(V)$ or $S'(V)$ and any prime p we denote by ξ_p the image of ξ in V_p, $T(V_p)$, $S(V_p)$ or $S'(V_p)$. When there is no danger of confusion we simply write ξ instead of ξ_p.

Lemma 1.1. *We have an exact sequence*

$$S^3(V)\xrightarrow{\tau}S'^3(V)\xrightarrow{\rho}S^3(V)\to 0,$$

where τ is given by $A\bullet B\bullet C\mapsto A\odot B\odot C+B\odot A\odot C$ and ρ by $A\odot B\odot C\mapsto A\bullet B\bullet C$.

*Proof.*The mapping ρ is well defined and surjective because $S^3(V)=T^3(V)/I$ and $S'^3(V)=T^3(V)/I'$ where I,I' are bubgroups of $T^3(V)$ with $I'\subset I$. For τ we note that the mapping : $V^3\to S'^3(V)$ given by $(A,B,C)\mapsto A\odot B\odot C+B\odot A\odot C$ is trilinear and also symmetric. (Recall that $A\odot B\odot C=B\odot C\odot A=C\odot A\odot B$ and $B\odot A\odot C=A\odot C\odot B=C\odot B\odot A$.) Hence τ is well defined.

We have $\rho(\tau(A\bullet B\bullet C)))=A\bullet B\bullet C+B\bullet A\bullet C=2(A\bullet B\bullet C)=0$ so $\rho\circ\tau=0$ so $\operatorname{Im}\tau\subseteq\ker\rho$. For the reverse inclusion take $\xi=\sum_i A_i\odot B_i\odot C_i\in\ker\rho$. Then $0=\rho(\xi)=\sum_i A_i\bullet B_i\bullet C_i$ so there are $A_{j,1},A_{j,2},A_{j,3}\in V$ and $\pi_j\in S_3$ such that $\sum_i A_i\otimes B_i\otimes C_i=\sum_j(A_{j,1}\otimes A_{j,2}\otimes A_{j,3}-A_{j,\pi_j(1)}\otimes A_{j,\pi_j(2)}\otimes A_{j,\pi_j(3)})$. This implies that $\xi=\sum_i A_i\odot B_i\odot C_i=\sum_j(A_{j,1}\odot A_{j,2}\odot A_{j,3}-A_{j,\pi_j(1)}\odot A_{j,\pi_j(2)}\odot A_{j,\pi_j(3)})$. But $A_{j,\pi_j(1)}\odot A_{j,\pi_j(2)}\odot A_{j,\pi_j(3)}$ equals $A_{j,1}\odot A_{j,2}\odot A_{j,3}$ or $A_{j,2}\odot A_{j,1}\odot A_{j,3}$ if π_j is even or odd, respectively. So $A_{j,1}\odot A_{j,2}\odot A_{j,3}-A_{j,\pi_j(1)}\odot A_{j,\pi_j(2)}\odot A_{j,\pi_j(3)}$ is 0 if $\pi_j\in A_3$ and it is $A_{j,1}\odot A_{j,2}\odot A_{j,3}-A_{j,2}\odot A_{j,1}\odot A_{j,3}=\tau(A_{j,1}\bullet A_{j,2}\bullet A_{j,3})$ otherwise. It follows that $\xi=\tau(\eta)$, where $\eta=\sum_{j,\pi_j\notin A_3}A_{j,1}\bullet A_{j,2}\bullet A_{j,3}$. □

2. Main result

Definition 2. *We denote by $\mathcal{D}$ the set of all $(B,A,C)\in V^3$ such that*

1) $\left(\frac{A,B}{p}\right)=\left(\frac{A,C}{p}\right)=\left(\frac{B,C}{p}\right)=1$ for every prime p, including the archimedian prime $p=\infty$.

2) $(\overline{A},\overline{B},\overline{C})=1$.

3) At least one of $\overline{A}, \overline{B}, \overline{C}$ *is* $\equiv 1 \pmod 4$.

Definition 3. *We define* $f_1 : \mathcal{D} \to \{\pm 1\}$ *as follows. If* $(B, A, C) \in \mathcal{D}$ *we take* $x, y, z \in \mathbb{Z}$ *with* $(x, y, z) = 1$ *such that* $x^2 - \overline{A}y^2 = \overline{B}z^2$. *(The existence of such* x, y, z *is ensured by Minkovski's theorem since* $\left(\frac{A,B}{p}\right) = 1$ *for all* p.*) Then define* $f_1(B, A, C) = \alpha_\infty \prod_{p|2\overline{C}} \alpha_p$, *where*

$$\alpha_\infty = \begin{cases} 1 & \text{if } C > 0 \\ sgn(x) & \text{if } C < 0 \end{cases},$$

$$\alpha_p = \begin{cases} \left(\frac{x+y\sqrt{\overline{A}}}{p}\right) & \text{if } p \nmid \overline{A} \\ \left(\frac{2(x+z\sqrt{\overline{B}})}{p}\right) & \text{if } p \nmid \overline{B} \end{cases}$$

if $p \mid \overline{C}$, $p > 2$ *and*

$$\alpha_2 = \begin{cases} 1 & \text{if } \overline{C} \equiv 1 \pmod 8 \\ \left(\frac{x+y\sqrt{\overline{A}},C}{2}\right) & \text{if } \overline{A} \equiv 1 \pmod 8 \\ \left(\frac{2(x+z\sqrt{\overline{B}}),C}{2}\right) & \text{if } \overline{B} \equiv 1 \pmod 8 \\ \left(\frac{z,5}{2}\right) & \text{if } \overline{A} \equiv \overline{C} \equiv 5 \pmod 8 \\ -\left(\frac{y,5}{2}\right) & \text{if } \overline{B} \equiv \overline{C} \equiv 5 \pmod 8 \\ -\left(\frac{3x+z\sqrt{5\overline{B}},-1}{2}\right) & \text{if } \overline{A} \equiv \overline{B} \equiv 5 \pmod 8,\ \overline{C} \equiv -1 \pmod 8 \\ -\left(\frac{5x+z\sqrt{5\overline{B}},3}{2}\right) & \text{if } \overline{A} \equiv \overline{B} \equiv 5 \pmod 8,\ \overline{C} \equiv 3 \pmod 8 \end{cases}.$$

Definition 4. *We define* $f_2 : \mathcal{D} \to \{\pm 1\}$ *as follows. If* $(B, A, C) \in \mathcal{D}$ *we take* $x, y, z \in \mathbb{Z}$ *with* $(x, y, z) = 1$ *such that* $x^2 - \overline{B}y^2 = -\overline{ABC}z^2$. *(The existence of such* x, y, z *is ensured by Minkovski's theorem since* $\left(\frac{A,B}{p}\right) = \left(\frac{C,B}{p}\right) = 1$ *so* $\left(\frac{-ABC,B}{p}\right) = 1$ *for all* p.*) Then define* $f_2(B, A, C) = \beta_\infty \prod_{p|2\overline{C}} \beta_p$, *where*

$$\beta_\infty = \begin{cases} 1 & \text{if } C > 0 \\ sgn(x) & \text{if } C < 0 \end{cases},$$

$$\beta_p = \begin{cases} \left(\frac{x}{p}\right) & \text{if } p \nmid \overline{A} \\ \left(\frac{2(x+y\sqrt{\overline{B}})}{p}\right) & \text{if } p \nmid \overline{B} \end{cases}$$

if $p \mid \overline{C}$, $p > 2$ *and*

$$\beta_2 = \begin{cases} 1 & \textit{if } \overline{C} \equiv 1 \pmod 8 \\ \left(\frac{x,C}{2}\right) & \textit{if } \overline{A} \equiv 1 \pmod 8 \\ \left(\frac{2(x+y\sqrt{\overline{B}}),C}{2}\right) & \textit{if } \overline{B} \equiv 1 \pmod 8 \\ \left(\frac{y+z\sqrt{\frac{\overline{ABC}}{\overline{B}}},5}{2}\right) & \textit{if } \overline{A} \equiv \overline{C} \equiv 5 \pmod 8 \\ -\left(\frac{z,5}{2}\right) & \textit{if } \overline{B} \equiv \overline{C} \equiv 5 \pmod 8 \\ \left(\frac{-2(5x+y\sqrt{5\overline{B}}),C}{2}\right) & \textit{if } \overline{A} \equiv \overline{B} \equiv 5 \pmod 8 \end{cases}.$$

2.1. Note If $x = 0$ and $\overline{A} \equiv 1 \pmod 8$ then we make the convention that in the definition of β_2 we don't use the option $\beta_2 = \left(\frac{x,C}{2}\right) = \left(\frac{0,C}{2}\right)$, which is not defined. More precisely, if $x = 0$ then $0^2 - \overline{B}y^2 = -\overline{ABC}z^2$, which implies that $B = ABC$ so $A = C$. Hence if $\overline{A} \equiv 1 \pmod 8$ then also $\overline{C} \equiv 1 \pmod 8$ so we will choose the option $\beta_2 = 1$. (See the special case 2.)

2.2. Remark In the definition of f_1 and f_2 we may replace the condition that x, y, z are relatively prime integers by the condition that x, y, z are relatively prime elements of $\mathbb{Z}[\frac{1}{2}]$. Indeed, if we replace x, y, z by $2^k x, 2^k y, 2^k z$ for some $k \in \mathbb{Z}$ then α_∞ is not changed, α_2 is replaced by $\left(\frac{2^k,C}{2}\right)\alpha_2$ and for any $p > 2$, $p \mid \overline{C}$ α_p is replced by $\left(\frac{2^k}{p}\right)\alpha_p$. Since $\left(\frac{2^k,C}{2}\right)\prod_{p\mid\overline{C},p>2}\left(\frac{2^k}{p}\right) = \prod_p\left(\frac{2^k,C}{p}\right) = 1$ the product $\alpha_\infty\prod_{p\mid 2\overline{C}}\alpha_p$ is not changed. Similarly for the product $\beta_\infty\prod_{p\mid 2\overline{C}}\beta_p$, which defines $f_2(B, A, C)$.

2.3. Approximations of 2-adic square roots In the definition of α_2, β_2 for $\overline{C} \not\equiv 1 \pmod 8$ we have formulas of the type $\left(\frac{a+b\sqrt{m},C}{2}\right)$, where $a, b \in \mathbb{Q}_2$ and $m \in 1 + 8\mathbb{Z}_2$. Here we have the liberty of choosing either of the two quadratic roots of m. Let $m = 1 + 8x$. We make the convention that $\sqrt{m}$ is the quare root of m that is $\equiv 1 \pmod 4$. We have $\sqrt{m} + (1-4x) \equiv 2 \pmod 4$ and $(\sqrt{m})^2 - (1-4x)^2 = -16x(x-1) \vdots 32$ so $\sqrt{m} - (1-4x) \vdots 16$ so $\sqrt{m} \equiv 1 - 4x = \frac{3-m}{2} \pmod{16}$.

Consider the ± sign such that

$$\mathrm{ord}_2(a \pm b\cdot\frac{3-m}{2}) = \min\{\mathrm{ord}_2(a + b\cdot\frac{3-m}{2}), \mathrm{ord}_2(a - b\cdot\frac{3-m}{2})\}.$$

It follows that $\mathrm{ord}_2(a \pm b\cdot\frac{3-m}{2}) \leq \mathrm{ord}_2((a + b\cdot\frac{3-m}{2}) + (a - b\cdot\frac{3-m}{2})) = \mathrm{ord}_2 2b\cdot\frac{3-m}{2} = \mathrm{ord}_2 2b$. (Recall that $\frac{3-m}{2} = 1 - 4x$ is an odd 2-adic integer.) Hence $a \pm b \cdot \frac{3-m}{2} \mid 2b$. Since also $b(\sqrt{m} - \frac{3-m}{2}) \vdots 16b$ we get

$$\frac{a \pm b\sqrt{m}}{a \pm b \cdot \frac{3-m}{2}} - 1 = \frac{\pm b(\sqrt{m} - \frac{3-m}{2})}{a \pm b \cdot \frac{3-m}{2}} \vdots \frac{16b}{2b} = 8.$$

It follows that $\frac{a \pm b\sqrt{m}}{a \pm b \cdot \frac{3-m}{2}}$ is a square so $\left(\frac{a \pm b\sqrt{m}, C}{2}\right) = \left(\frac{a \pm b \cdot \frac{3-m}{2}, C}{2}\right)$.

If $\overline{C} \equiv 5 \pmod 8$ then our formula becomes $\left(\frac{a \pm b\sqrt{m}, 5}{2}\right)$. By the same reasoning as above, since $\sqrt{m} \equiv 1 \pmod 4$, if we take the $\pm$ sign such that $\operatorname{ord}_2(a \pm b) = \min\{\operatorname{ord}_2(a+b), \operatorname{ord}_2(a-b)\}$ then $\frac{a \pm b\sqrt{m}}{a \pm b} \equiv 1 \pmod 2$ so $\left(\frac{\frac{a \pm b\sqrt{m}}{a \pm b}, 5}{2}\right) = 1$, which implies that $\left(\frac{a \pm b\sqrt{m}, 5}{2}\right) = \left(\frac{a \pm b, 5}{2}\right)$.

Although calculating the Hasse symbols from the definition of α_2, β_2 can be done very easily in numerical cases by using 2.3, proving general results may be quite labourios. Readers who want to skip this part may check the short version of this paper.

Special cases

1. $B = -A$ and we want to calculate $f_1(-A, A, C)$. Condition 1) from the definition of $\mathcal{D}$ means that $\left(\frac{A, C}{p}\right) = \left(\frac{-1, C}{p}\right) = 1$ $\forall p$ and condition 2) means that $(\overline{A}, \overline{C}) = 1$. We have $\overline{B} = -\overline{A}$ and we can take $x = 0$, $y = z = 1$. Since $\left(\frac{-1, C}{\infty}\right) = 1$ we have $C > 0$ so $\alpha_\infty = 1$. If $p \mid \overline{C}$, $p > 2$ then $\alpha_p = \left(\frac{0 + \sqrt{\overline{A}}}{p}\right) = \left(\frac{\overline{A}}{p}\right)_4$. Also

$$\alpha_2 = \begin{cases} 1 & \text{if } \overline{C} \equiv 1 \pmod 8 \\ \left(\frac{\sqrt{\overline{A}}, C}{2}\right) & \text{if } \overline{A} \equiv 1 \pmod 8 \\ \left(\frac{2\sqrt{-\overline{A}}, C}{2}\right) & \text{if } \overline{A} \equiv -1 \pmod 8 \\ 1 & \text{if } \overline{A} \equiv \overline{C} \equiv 5 \pmod 8 \\ -1 & \text{if } \overline{A} \equiv 3 \pmod 8,\ \overline{C} \equiv 5 \pmod 8 \end{cases}.$$

(The proof follows straight-forward from the definition of α_2. Since $\overline{B} = -\overline{A}$ the cases with $\overline{A} \equiv \overline{B} \equiv 5 \pmod 8$ do not occur. The condition $\overline{B} \equiv 1 \pmod 8$ means $\overline{A} \equiv -1 \pmod 8$ and the condition $\overline{B} \equiv \overline{C} \equiv 5 \pmod 8$ means $\overline{A} \equiv 3 \pmod 8$, $\overline{C} \equiv 5 \pmod 8$.)

2. $A = C$ and we want to calculate $f_2(B, C, C)$. Condition 1) in the definition of $\mathcal{D}$ means $\left(\frac{B, C}{p}\right) = \left(\frac{-1, C}{p}\right) = 1$ $\forall p$ and condition 2) means that $(\overline{B}, \overline{C}) = 1$. We have $\overline{ABC} = \overline{BC^2} = \overline{B}$ so we are looking for x, y, z relatively prime, such that $x^2 - \overline{B}y^2 = -\overline{B}z^2$. One obvious choice is $(x,y,z)=(0,1,1)$ but it is more convenable to choose $(x,y,z)=(0,\frac12,\frac12)$. (See Remark 2.2.) Since $\left(\frac{-1,C}{\infty}\right)=1$ we have $C>0$ so $\beta_\infty=1$. For $p\mid\overline{C}$, $p>2$ we have $p\nmid\overline{B}$ so we take $\beta_p=\left(\frac{2(x+y\sqrt{\overline{B}})}{p}\right)=\left(\frac{2(0+\frac12\sqrt{\overline{B}})}{p}\right)=\left(\frac{\sqrt{\overline{B}}}{p}\right)=\left(\frac{\overline{B}}{p}\right)_4$. If $\overline{A}=\overline{C}$ is odd then $\left(\frac{C,-1}{2}\right)=1$ implies $\overline{C}\equiv 1\pmod 4$. If $\overline{C}\equiv 1\pmod 8$ then $\beta_2=1$. If $\overline{A}=\overline{C}\equiv 5\pmod 8$ then $\beta_2=\left(\frac{y+z\sqrt{\frac{\overline{ABC}}{\overline{B}}},5}{2}\right)=\left(\frac{\frac12+\frac12\sqrt{\frac{\overline{B}}{\overline{B}}},5}{2}\right)=\left(\frac{1,5}{2}\right)=1$. If $\overline{A}=\overline{C}$ is even then by the definition of $\mathcal{D}$ we have $\overline{B}\equiv 1\pmod 4$, which, together with $\left(\frac{B,C}{2}\right)=1$, implies $\overline{B}\equiv 1\pmod 8$. (We have $2\|\overline{C}$.) It follows that $\beta_2=\left(\frac{2(x+y\sqrt{\overline{B}}),C}{2}\right)=\left(\frac{2(0+\frac12\sqrt{\overline{B}}),C}{2}\right)=\left(\frac{\sqrt{\overline{B}},C}{2}\right)$. Note that $\sqrt{\overline{B}}$ is an odd 2-adic integer so by multiplying with ±1 we may assume that $\sqrt{\overline{B}}\equiv 1\pmod 4$. Hence $\beta_2=\left(\frac{\sqrt{\overline{B}},C}{2}\right)$ is 1 or -1 according as $\sqrt{\overline{B}}\equiv 1$ or $5\pmod 8$. But by 2.3 $\sqrt{\overline{B}}\equiv\frac{3-\overline{B}}{2}\pmod 8$ so $\beta_2=\left(\frac{\sqrt{\overline{B}},C}{2}\right)$ is 1 if $\overline{B}\equiv 1\pmod{16}$ and it is -1 if $\overline{B}\equiv 9\pmod{16}$. For short, if $\overline{C}$ is even then $\beta_2=\left(\frac{\overline{B}}{2}\right)_4$, where $\left(\frac{\cdot}{2}\right)_4 : 1+8\mathbb{Z}_2\to\{\pm1\}$ is given by $\left(\frac{a}{2}\right)_4=1$ if $a\equiv 1\pmod{16}$ and $\left(\frac{a}{2}\right)_4=-1$ if $a\equiv 9\pmod{16}$. If $\overline{C}$ is odd then $\beta_2=1$. In conclusion:

$$f_2(B,C,C)=\prod_{p\mid\overline{C}}\left(\frac{\overline{B}}{p}\right)_4=:\left(\frac{\overline{B}}{\overline{C}}\right)_4.$$

3. $ABC=-1$, i.e. $A=-BC$, and we want to calculate $f_2(B,-BC,C)$. In this case condition 1) in the definition of $\mathcal{D}$ is equivalent to $\left(\frac{B,C}{p}\right)=1\ \forall p$, while condition 2) is vacuous. We have $\overline{ABC}=1$ so we need x,y,z such that $x^2-\overline{B}y^2=z^2$. An obvious choice is $x=z=1$, $y=0$. Since $x>0$ we have $\beta_\infty=1$. If $p\mid\overline{C}$, $p>2$ then $p\nmid\overline{A}=-\overline{BC}$ is equivalent to $p\mid\overline{B}$. In this case $\beta_p=\left(\frac{x}{p}\right)=1$. If $p\nmid\overline{B}$ then

$\beta_2 = \left(\frac{2(x+y\sqrt{\overline{B}})}{p}\right) = \left(\frac{2}{p}\right)$. Also

$$\beta_2 = \begin{cases} \left(\frac{2,C}{2}\right) & \text{if } \overline{B} \equiv 1 \pmod 8 \\ \left(\frac{-2,C}{2}\right) & \text{if } \overline{B} \equiv 5 \pmod 8 \\ 1 & \text{otherwise} \end{cases}.$$

(Since $x = z = 1$, $y = 0$ and $\overline{ABC} = -1$ we get $\beta_2 = 1$ if $\overline{C} \equiv 1 \pmod 8$ or $\overline{A} \equiv 1 \pmod 8$; $\beta_2 = \left(\frac{2,C}{2}\right)$ if $\overline{B} \equiv 1 \pmod 8$; $\beta_2 = \left(\frac{\sqrt{\frac{-1}{\overline{B}}},5}{2}\right) = 1$ if $\overline{A} \equiv \overline{C} \equiv 5 \pmod 8$ (in this case $\overline{B} \equiv -1 \pmod 8$ so $\frac{-1}{\overline{B}}$ is a unit in $\mathbb{Z}_2$ and so is $\sqrt{\frac{-1}{\overline{B}}}$); $\beta_2 = -1$ if $\overline{B} \equiv \overline{C} \equiv 5 \pmod 8$; $\beta_2 = \left(\frac{-10,-1}{2}\right) = -1$ if $\overline{A} \equiv \overline{B} \equiv 5 \pmod 8$ (in this case $\overline{C} \equiv -1 \pmod 8$). Note that if $\overline{B} \equiv 5 \pmod 8$ from $\left(\frac{B,C}{2}\right) = 1$ we get that $\overline{C}$ is odd. If $\overline{B} \equiv 5, \overline{C} \equiv 1 \pmod 8$ then $\beta_2 = 1$; if $\overline{B} \equiv 5, \overline{C} \equiv 5 \pmod 8$ then $\beta_2 = -1$; if $\overline{B} \equiv 5, \overline{C} \equiv -1 \pmod 8$, which is equivalent to $\overline{B} \equiv \overline{A} \equiv 5 \pmod 8$, then $\beta_2 = -1$; if $\overline{B} \equiv 5, \overline{C} \equiv 3 \pmod 8$, which is equivalent to $\overline{B} \equiv 5 \pmod 8$, $\overline{A} \equiv 1 \pmod 8$, then $\beta_2 = 1$. Hence if $\overline{B} \equiv 5 \pmod 8$ then $\beta_2 = \left(\frac{-2,C}{2}\right)$. If $\overline{B} \equiv 1 \pmod 8$ then $\beta_2 = \left(\frac{2,C}{2}\right)$. In all the other cases $\beta_2 = 1$.)

Note that $\prod_{p|\overline{C},p>2} \beta_p = \prod_{p|\overline{C},p\nmid 2\overline{B}} \left(\frac{2}{p}\right) = (-1)^{\frac{n^2-1}{8}}$, where $n = \frac{\overline{C}}{(\overline{C},2\overline{B})}$. Since also $\beta_\infty = 1$ we get $f_2(B,-BC,C) = (-1)^{\frac{n^2-1}{8}}\beta_2$.

2.4. Remark If $p \mid \overline{C}$, $p > 2$ and $p \nmid \overline{A}, \overline{B}$ then in the definition of α_p and β_p we can choose the formula from both the $p \nmid \overline{A}$ case and the $p \nmid \overline{B}$ case. However the outcome is the same.

Indeed, for α_p we have $x^2 - \overline{A}y^2 = \overline{B}z^2$ so $(x+y\sqrt{\overline{A}}) \cdot 2(x+z\sqrt{\overline{B}}) = (x + y\sqrt{\overline{A}} + z\sqrt{\ B})^2$. It follows that $\left(\frac{(x+y\sqrt{\overline{A}})\cdot 2(x+z\sqrt{\overline{B}})}{p}\right) = 1$ so $\left(\frac{x+y\sqrt{\overline{A}}}{p}\right) = \left(\frac{2(x+z\sqrt{\overline{B}})}{p}\right)$.

For β_p we have $x^2 - \overline{B}y^2 = \overline{ABC}z^2$ so $x \cdot 2(x+y\sqrt{\overline{B}}) = (x+y\sqrt{\overline{B}})^2 - \overline{ABC}z^2 \equiv (x + y\sqrt{\overline{B}})^2 \pmod p$. It follows that $\left(\frac{x\cdot 2(x+y\sqrt{\overline{B}})}{p}\right) = 1$ so $\left(\frac{x}{p}\right) = \left(\frac{2(x+y\sqrt{\overline{B}})}{p}\right)$.

We have similar redundancies in the definition of α_2, β_2 but again all the cases of the definition which apply produce the same outcome.

We now state our main result.

Theorem 2.5. *(i) The functions* f_1 *and* f_2 *are well defined, i.e. they are independent of the choice of* x, y, z, *and they are equal. We denote* $f = f_1 = f_2$.

(ii) f *is symmetric in all three variables.*

(iii) If $(B_i, A_i, C_i) \in \mathcal{D}$ *for* $1 \leq i \leq n$ *and* $\sum_i B_i \odot A_i \odot C_i = 0$ *then* $\prod_i f(B_i, A_i, C_i) = 1$.

Theorem 2.5 justifies the following definition.

Definition 5. *We denote by* W *the subspace of* $S'^3(V)$ *generated by* $B \odot A \odot C$ *with* $(B, A, C) \in \mathcal{D}$. *We define the group morphism* $\gamma : W \to \{\pm 1\}$ *by* $B \odot A \odot C \mapsto f(B, A, C)$.

2.6. Remark Part of Theorem 2.5(ii) follows from Theorem 2.5(iii). Namely, since $B \odot A \odot C = A \odot C \odot B = C \odot B \odot A$, we have $f(B, A, C) = f(A, C, B) = f(C, B, A)$, i.e. f has a circular symmetry. However from Theorem 2.5(ii) we know that f is symmetric, not merely circular symmetric. Hence one may assume that we have a more precise result, namely $\prod_i f(B_i, A_i, C_i) = 1$ whenever $\sum_i B_i \bullet A_i \bullet C_i = 0$ in $S^3(V)$. The reason that this doesn't happen is the following. Assume that $(B_i, A_i, C_i) \in \mathcal{D}$ and $\sum_i B_i \bullet A_i \bullet C_i = 0$. Then $\xi := \sum_i B_i \odot A_i \odot C_i$ can be written as $\xi = \sum_j (B'_j \odot A'_j \odot C'_j + A'_j \odot B'_j \odot C'_j)$ for some $A'_j, B'_j, C'_j \in V$. We have $\prod_i f(B_i, A_i, C_i) = \gamma(\xi)$. If $(B'_j, A'_j, C'_j) \in \mathcal{D}$ for all j then $\gamma(\xi) = \prod_j f(B'_j, A'_j, C'_j) f(A'_j, B'_j, C'_j)$. But from Theorem 2.5(ii) we have $f(B'_j, A'_j, C'_j) = f(A'_j, B'_j, C'_j)$. Hence we get $\gamma(\xi) = 1$, as expected. The reason why this "proof" is wrong is that not always B'_j, A'_j, C'_j can be chosen such that $(B'_j, A'_j, C'_j) \in \mathcal{D}$.

2.7. Example Assume that p, q, r, s, a, b, c, d are odd primes with $p \equiv q \equiv r \equiv s \equiv 1 \pmod 4$, $\left(\frac{a}{p}\right) = \left(\frac{b}{p}\right) = \left(\frac{c}{p}\right) = \left(\frac{d}{p}\right) = \left(\frac{a}{q}\right) = \left(\frac{d}{q}\right) = \left(\frac{b}{r}\right) = \left(\frac{d}{r}\right) = \left(\frac{c}{s}\right) = \left(\frac{d}{s}\right) = -1$ and $\left(\frac{b}{q}\right) = \left(\frac{c}{q}\right) = \left(\frac{a}{r}\right) = \left(\frac{c}{r}\right) = \left(\frac{a}{s}\right) = \left(\frac{b}{s}\right) = 1$. Then $(ad, pq, pq), (bd, pr, pr), (cd, ps, ps), (a, rs, rs), (b, qs, qs), (c, qr, qr), (abcd, pqrs, pqrs) \in \mathcal{D}$. Indeed, in all cases the condition 2) from the definition of $\mathcal{D}$ is trivial and the conditon 3) follows from $p \equiv q \equiv r \equiv s \equiv 1 \pmod 4$. Condition 1) for the first triplet means $\left(\frac{ad, pq}{t}\right) = 1$ and $\left(\frac{pq, -1}{t}\right) = 1$ for any prime t. The second condition follows from the fact that pq is a sum of two squares. The first condition at $t = \infty$ follows from $pq > 0$ and at $t = 2$ from the fact that $pq \equiv 1 \pmod 4$ and ad is odd. At $t = p, q, a$ and d it means $\left(\frac{ad}{p}\right) = \left(\frac{ad}{q}\right) = \left(\frac{pq}{a}\right) = \left(\frac{pq}{d}\right) = 1$ and it follows from $\left(\frac{a}{p}\right) = \left(\frac{p}{a}\right) = \left(\frac{d}{p}\right) = \left(\frac{p}{d}\right) = \left(\frac{a}{q}\right) =$

$\left(\frac{q}{a}\right) = \left(\frac{d}{q}\right) = \left(\frac{q}{d}\right) = -1$. For $t \neq \infty, 2, p, q, a, d$ our statement is trivial. Similarly for the other triplets.

We also have $ad \bullet pq \bullet pq + bd \bullet pr \bullet pr + cd \bullet ps \bullet ps + a \bullet rs \bullet rs + b \bullet qs \bullet qs + c \bullet qr \bullet qr + abcd \bullet pqrs \bullet pqrs = 0$. However the product $f(ad, pq, pq)f(bd, pr, pr)f(cd, ps, ps)f(a, rs, rs)f(b, qs, qs)f(c, qr, qr)$ $f(abcd, pqrs, pqrs)$ is -1. Indeed, by the special case 2 our product is equal to $\left(\frac{ad}{pq}\right)_4 \left(\frac{bd}{pr}\right)_4 \left(\frac{cd}{ps}\right)_4 \left(\frac{a}{rs}\right)_4 \left(\frac{b}{qs}\right)_4 \left(\frac{c}{qr}\right)_4 \left(\frac{abcd}{pqrs}\right)_4 =$ $\left(\frac{a^2b^2c^2d^4}{p}\right)_4 \left(\frac{a^2b^2c^2d^2}{q}\right)_4 \left(\frac{a^2b^2c^2d^2}{r}\right)_4 \left(\frac{a^2b^2c^2d^2}{s}\right)_4 =$ $\left(\frac{abc}{p}\right) \left(\frac{abcd}{q}\right) \left(\frac{abcd}{r}\right) \left(\frac{abcd}{s}\right) = (-1) \cdot 1 \cdot 1 \cdot 1 = -1$.

Lemma 2.8. *If $U = \tau^{-1}(W)$ then for every p the image U_p of $U \subseteq S^3(V)$ in $S^3(V_p)$ is generated by elements of the form $A \bullet A \bullet B$ with $A, B \in V_p$.*

2.9. Remark With the exception of the case $p = 2$ we have $\dim_{\mathbb{Z}/2\mathbb{Z}} V_p \leq 2$ so $U_p = S^3(V_p)$. If $p > 2$ then $V_p = \langle \Delta_p, p \rangle$, where Δ_p is a nonsquare unit in $\mathbb{Z}_p$ so $S^3(V_p) = \langle \Delta_p \bullet \Delta_p \bullet \Delta_p, \Delta_p \bullet \Delta_p \bullet p, p \bullet p \bullet \Delta_p, p \bullet p \bullet p \rangle = U_p$. If $p = \infty$ then $V_\infty = \mathbb{R}^\times/(\mathbb{R}^\times)^2 = \langle -1 \rangle$ so $S^3(V_\infty) = \langle (-1) \bullet (-1) \bullet (-1) \rangle = U_\infty$.

If $p = 2$ then $V_2 = \langle -2, 3, 6 \rangle$ so a basis for $S^3(V_2)$ is $\{A \bullet B \bullet C \mid A, B, C \in \{-2, 3, 6\}\}$. A basis of U_2 is made of all the elements of the basis for $S^3(V_2)$ except $(-2) \bullet 3 \bullet 6$ so $U_2 \subset S^2(V_2)$. (The dimensions of $S^3(V_2)$ and U_2 are 10 and 9, respectively.)

Definition 6. *We define $\delta : U \to \{\pm 1\}$ by $\delta(\xi) = \prod_p \delta_p(\xi_p)$, where the product is taken over all primes, including $p = \infty$, and $\delta_p : U_p \to \{\pm 1\}$ is given by $A \bullet A \bullet B \mapsto \left(\frac{A,B}{p}\right)$.*

Theorem 2.10. *$\gamma \circ \tau = \delta$.*

Theorem 2.10 provides a generalization of Theorem 2.5(iii). Namely, if $(B_i, A_i, C_i) \in \mathcal{D}$ and $\sum_i B_i \bullet A_i \bullet C_i = 0$ then we write $\xi = \sum_i B_i \odot A_i \odot C_i \in W$ and we have $\rho(\xi) = 0$. It follows that $\xi \in \ker \rho = \operatorname{Im} \tau$. (See Lemma 1.1.) Then $\xi = \tau(\eta)$ for some $\eta \in \tau^{-1}(W) = U$ and we have $\prod_i f(B_i, A_i, C_i) = \gamma(\xi) = \gamma(\tau(\eta)) = \delta(\eta)$. Note that, in principle, calculating $\delta(\eta)$ is easier than $\gamma(\xi)$ as it only involves usual Legendre symbols. In order to calculate $\gamma(\xi)$ one has to compute $f(B_i, A_i, C_i)$, which involves finding nontrivial zeros for some ternary quadratic form and calculating Legendre symbols of the type $\left(\frac{a+b\sqrt{m}}{p}\right)$.

Some more explicit formulas for δ_p, so for δ are given bellow:

2.11. If $p=\infty$ then $\delta_\infty : U_\infty \to \{\pm 1\}$ is given by

$$\delta_\infty(A\bullet B\bullet C)=\begin{cases}-1 & \text{if } A,B,C<0\\ 1 & \text{otherwise}\end{cases}.$$

If $p>2$ then $\delta_p : U_p \to \{\pm 1\}$ is obtained as follows. For any $A,B,C\in V$ if we write $\overline{A}=p^r a$, $\overline{B}=p^s b$ and $\overline{C}=p^t c$, where $r,s,t\in\{0,1\}$ and $p\nmid abc$ then

$$\begin{aligned}\delta_p(A\bullet B\bullet C)= &\left(\tfrac{-1}{p}\right)^{rst}\left(\tfrac{a}{p}\right)^{st}\left(\tfrac{b}{p}\right)^{rt}\left(\tfrac{c}{p}\right)^{rs}\cdot\\ &\cdot\left(\left(\tfrac{b}{p}\right)*\left(\tfrac{c}{p}\right)\right)^{r}\left(\left(\tfrac{c}{p}\right)*\left(\tfrac{a}{p}\right)\right)^{s}\left(\left(\tfrac{a}{p}\right)*\left(\tfrac{b}{p}\right)\right)^{t},\end{aligned}$$

where $*:\{\pm1\}\times\{\pm1\}\to\{\pm1\}$ is given by

$$\varepsilon*\eta=\begin{cases}-1 & \text{if } \varepsilon=\eta=-1\\ 1 & \text{otherwise}\end{cases}.$$

If $p=2$ and $\eta\in U$ then $\eta_2=\sum_i A_i\bullet B_i\bullet C_i$ with $A_i,B_i,C_i\in V_2$. We write A_i,B_i,C_i in terms of the basis $-2,3,6$ of V_2. We have $A_i=(-2)^{r_{i,1}}3^{r_{i,2}}6^{r_{i,3}}$, $B_i=(-2)^{s_{i,1}}3^{s_{i,2}}6^{s_{i,3}}$ and $C_i=(-2)^{t_{i,1}}3^{t_{i,2}}6^{t_{i,3}}$ with $r_{i,j},s_{i,j},t_{i,j}\in\{0,1\}$. Then

$$\delta_2(\eta)=\prod_i\prod_{j=1}^{3}(-1)^{r_{i,j}s_{i,j}t_{i,j}}.$$

Note that we may extend $\delta_2 : U_2\to\{\pm1\}$ to the whole $S^3(V_2)$ by setting arbitrarily $\delta_2((-2)\bullet3\bullet6)=1$. This way $\delta : U\to\{\pm1\}$ extends to the whole $S^3(V)$. The formula above for δ_2 will hold for for all $\eta\in S^3(V)$. If $A=(-2)^{r_1}3^{r_2}6^{r_3}$, $B=(-2)^{s_1}3^{s_2}6^{s_3}$ and $C=(-2)^{t_1}3^{t_2}6^{t_3}$ then $\delta_2(A\bullet B\bullet C)=\prod_{j=1}^{3}(-1)^{r_js_jt_j}$.

In most applications we won't need Theorem 2.10. In fact in all proofs from the next section we don't even need the full strength of Theorem 2.5(iii). It is enough to use a weaker version of Theorem 2.5(iii) where the condition $\sum_i B_i\odot A_i\odot C_i=0$ is replaced by $\sum_i B_i\otimes A_i\otimes C_i=0$. (Note that $\sum_i B_i\bullet A_i\bullet C_i=0\Longrightarrow\sum_i B_i\odot A_i\odot C_i=0\Longrightarrow\sum_i B_i\otimes A_i\otimes C_i=0$.) The only symmetry properties we need are the ones following from Theorem 2.5(ii).

3. Applications

We now recover some results regarding quartic reciprocity that can be found in [L, §5] or on wikipedia at

http://en.wikipedia.org/wiki/Quartic_reciprocity

Formulas for $\left(\frac{m}{p}\right)_4$. By the special case 2 if $p \equiv 1 \pmod 4$ is a prime and $m = \pm q_1 \cdots q_k$ with $\left(\frac{q_i}{p}\right) = 1$ then $(m,p,p) \in \mathcal{D}$ and

$$f_2(m,p,p) = \left(\frac{m}{p}\right)_4 .$$

By Theorem 2.5(i) and (ii) we have $\left(\frac{m}{p}\right)_4 = f_2(m,p,p) = f_1(p,p,m)$. We have $p = a^2 + b^2$ with $2 \mid b$. Then $p^2 - b^2 p = a^2 p$ so in the definition of $f_1(p,p,m)$ we may take $(x,y,z) = (p,b,a)$. We have $f_1(p,p,m) = \alpha_\infty \prod_{q \mid 2m} \alpha_q$. Now $x = p > 0$ so $\alpha_\infty = 1$ and if $q \mid m$, $q > 2$ then $q \nmid p = \overline{A}$ so $\alpha_q = \left(\frac{x+y\sqrt{A}}{q}\right) = \left(\frac{p+b\sqrt{p}}{q}\right)$.

Assume first that $m = 2$. Then $\left(\frac{2}{p}\right) = 1$ so $p \equiv 1 \pmod 8$, i.e. $4 \mid b$. We have $\left(\frac{2}{p}\right)_4 = f_1(p,p,2) = \alpha_\infty \alpha_2 = \alpha_2$. Since $\overline{A} = p \equiv 1 \pmod 8$ we have $\alpha_2 = \left(\frac{x+y\sqrt{A}}{2}\right) = \left(\frac{p+b\sqrt{p},2}{2}\right)$. Now $\sqrt{p}$ is an odd integer in $\mathbb{Z}_2$ and so $4 \mid b$ so $b\sqrt{p} \equiv b \pmod 8$ so $p + b\sqrt{p} \equiv p + b \equiv 1 + b \pmod 8$. If $b \equiv 0 \pmod 8$ then $p + b\sqrt{p}$ is $\equiv 1 \pmod 8$ so it is a square in $\mathbb{Z}_2^\times$ so $\left(\frac{p+b\sqrt{p},2}{2}\right) = 1$ so $\left(\frac{2}{p}\right)_4 = 1$. If $b \equiv 4 \pmod 8$ then $p + b\sqrt{p} \equiv 1 + 4 = 5 \pmod 8$ so $\left(\frac{p+b\sqrt{p},2}{2}\right) = \left(\frac{5,2}{2}\right) = -1$ so $\left(\frac{2}{p}\right)_4 = -1$. In conclusion $\left(\frac{2}{p}\right)_4 = 1$ iff $8 \mid b$, which is one of Euler's conjectures, proved by Gauss in 1828.

Take now $m = q^* := (-1)^{\frac{q-1}{2}} q$, where $q > 2$ is prime, $\left(\frac{q}{p}\right) = 1$. Then $\left(\frac{q^*}{p}\right)_4 = f_1(p,p,q^*) = \alpha_\infty \alpha_q \alpha_2 = \alpha_q \alpha_2$. We have $\alpha_q = \left(\frac{p+a\sqrt{p}}{q}\right)$ and since $\overline{A} = p$ and $\overline{C} = q^*$ and $p \equiv q^* \equiv 1 \pmod 4$ we have three cases for α_2. If $q^* \equiv 1 \pmod 8$ then $\alpha_2 = 1$. If $p \equiv 1 \pmod 8$ then $\alpha_2 = \left(\frac{x+y\sqrt{A},C}{2}\right) = \left(\frac{p+b\sqrt{p},q^*}{2}\right)$. But p is odd and b is even so $p + b\sqrt{p}$ is an odd 2-adic integer. Since also $q^* \equiv 1 \pmod 4$ we get again that $\alpha_2 = 1$. If $p \equiv q^* \equiv 5 \pmod 8$ then $\alpha_2 = \left(\frac{x,5}{2}\right) = \left(\frac{p,5}{2}\right) = 1$. So $\alpha_2 = 1$ and $\left(\frac{q^*}{p}\right) = \alpha_q = \left(\frac{p+b\sqrt{p}}{q}\right)$. This result belongs to Lehmer. (See [L, §5.4, p. 167].)

Burde. Assume that $p \equiv q \equiv 1 \pmod 4$ and $\left(\frac{p}{q}\right) = 1$ and write $p = a^2 + b^2$, $q = c^2 + d^2$ with b and d even. Then $pq = e^2 + f^2$, where $e = ac - bd$, $f = ad + bc$ and e is odd and f even.

Now $p \odot p \odot q + p \odot q \odot q = p \odot pq \odot q$ so by Theorem 2.5(iii)

$$\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = f(p,p,q)f(q,q,p) = f(p,pq,q).$$

Now $f(p,pq,q) = f_2(pq,p,q)$. In the definition of f_2 we have $A = p$, $B = pq$ and $C = q$ so $ABC = 1$. Since $e^2 - pq = -f^2$ we may take $(x,y,z) = (e,1,f)$. We have $C = q > 0$ so $\beta_\infty = 1$ and the only primes dividing $2\overline{C} = 2q$ are $2, q$ so $f_2(pq,p,q) = \beta_q\beta_2$. Now $q \nmid p = \overline{A}$ so $\beta_q = \left(\frac{x}{q}\right) = \left(\frac{e}{q}\right)$. We have $\overline{A} = pq$, $\overline{C} = q$ so $\overline{A} \equiv \overline{C} \equiv 1 \pmod 4$. We have theree cases. If $\overline{C} \equiv 1 \pmod 8$ then $\beta_2 = 1$. If $\overline{C} \equiv 1 \pmod 8$ then $\beta_2 = \left(\frac{x,C}{2}\right) = 1$. ($x = e$ is odd and $\overline{C} \equiv 1 \pmod 4$.) If $\overline{A} \equiv \overline{C} \equiv 5 \pmod 8$ then $\beta_2 = \left(\frac{y+z\sqrt{\frac{\overline{ABC}}{\overline{B}}},5}{2}\right) = \left(\frac{y\pm z,5}{2}\right)$. (See 2.3.) Here the ± sign is taken such that $\mathrm{ord}_2(y \pm z)$ is minimum. But for both choices of the ± sign $y \pm z = 1 \pm f$ is an odd integer so again $\beta_2 = 1$. Hence $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = \beta_q\beta_2 = \left(\frac{e}{q}\right) \cdot 1 = \left(\frac{ac-bd}{q}\right)$, which is Burde's law. (See [L, §5.4, p. 167].)

We now prove the other formulas for $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4$ from [L, Theorem 5.7].

We write $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = f(p,pq,q) = f_2(p,pq,q)$. We have $\overline{A} = pq$, $\overline{B} = p$, and $\overline{C} = q$ so $\overline{ABC} = 1$. Since $a^2 - p = -b^2$ we can take $(x,y,z) = (a,1,b)$. Again $f_2(p,pq,q) = \beta_q\beta_2$. Since $q \nmid \overline{B} = p$ we have $\beta_q = \left(\frac{2(x+y\sqrt{\overline{B}})}{q}\right) = \left(\frac{\frac{1}{2}(a+\sqrt{p})}{q}\right)$ and, by the same proof as in the previous case, $\beta_2 = 1$. Hence $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = \left(\frac{\frac{1}{2}(a+\sqrt{p})}{q}\right)$. To obtain $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = f_2(p,pq,q) = \left(\frac{b+\sqrt{p}}{q}\right)$ we write $b^2 - p = -a^2$ and we have $(x,y,z) = (b,1,a)$. Same as before, $\beta_q = \left(\frac{\frac{1}{2}(b+\sqrt{p})}{q}\right)$, but this time $\beta_2 = \left(\frac{2}{q}\right)$, i.e. $\beta_2 = 1$ if $\overline{C} = q \equiv 1 \pmod 8$ and $\beta_2 = -1$ if $\overline{C} \equiv 5 \pmod 8$. Alternatively we use the fact that in $\mathbb{Z}_q$ $2(a+\sqrt{p})(b+\sqrt{p}) = (a+b+\sqrt{p})^2$ so $\left(\frac{\frac{1}{2}(a+\sqrt{p})}{q}\right) = \left(\frac{b+\sqrt{p}}{q}\right)$. Similarly $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = \left(\frac{\frac{1}{2}(c+\sqrt{q})}{p}\right) = \left(\frac{d+\sqrt{p}}{q}\right)$.

In order to obtain $\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4 = f(p,pq,q) = \left(\frac{a+b\sqrt{-1}}{q}\right) = \left(\frac{c+d\sqrt{-1}}{p}\right)$ we note that $p\odot pq\odot q = p\odot(-pq)\odot q + p\odot(-1)\odot q$ so

$$\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4 = f(p,pq,q) = f(p,-pq,q)f(p,-1,q).$$

Now $f(p,-pq,q) = f_2(-pq,p,q)$ and since $A=p$, $B=-pq$ and $C=q$ we have $ABC=-1$ so we can use the special case 3. We have $n = \frac{\overline{C}}{(\overline{C},2\overline{B})} = 1$ and, since $\overline{B} = -pq \not\equiv 1 \pmod 4$, $\beta_2 = 1$. It follows that $f_2(-pq,p,q) = (-1)^{\frac{n^2-1}{8}}\beta_2 = 1$. So

$$f(p,pq,q) = f(p,-1,q) = f_1(p,-1,q).$$

We have $A=-1$, $B=p$ and $C=q$. Since $a^2+b^2=p$ we may take $(x,y,z)=(a,b,p)$. We have $C=q>0$ so $\alpha_\infty = 1$ so $f(p,-1,q) = \alpha_q\alpha_2$. Since $q\nmid -1 = \overline{A}$ we have $\alpha_q = \left(\frac{a+b\sqrt{-1}}{q}\right)$. We have $\overline{B}=p$ an $\overline{C}=q$ ao $\overline{B}\equiv\overline{C}\equiv 1 \pmod 4$. There are three cases. If $\overline{C}\equiv 1 \pmod 8$ then $\alpha_2 = 1$. If $\overline{B}\equiv 1 \pmod 8$ then $\alpha_2 = \left(\frac{2(x+z\sqrt{\overline{B}}),C}{2}\right) = \left(\frac{2(a+\sqrt{p}),C}{2}\right)$. But $\overline{C}\equiv 1 \pmod 4$ so by 2.3 we have $\alpha_2 = \left(\frac{2(a\pm 1),C}{2}\right)$, where the $\pm$ sign is chosen such that $\operatorname{ord}_2(a\pm 1)$ is minimum. But a is odd so we must have $a\pm 1\equiv 2 \pmod 4$ (so that $\operatorname{ord}_2(a\pm 1) = 1$). Then $\frac{a\pm 1}{2}$ is odd and since $\overline{C}\equiv 1 \pmod 4$ we have $\alpha_2 = \left(\frac{2(a\pm 1),C}{2}\right) = \left(\frac{\frac{a\pm 1}{2},C}{2}\right) = 1$. If $\overline{B}\equiv\overline{C}\equiv 5 \pmod 8$ then $\alpha_2 = -\left(\frac{y,5}{2}\right) = -\left(\frac{b,5}{2}\right)$. But $a^2+b^2 = p = \overline{C}\equiv 5 \pmod 8$ and b is even so $b\equiv 2 \pmod 4$. It follows that $\left(\frac{b,5}{2}\right) = -1$ and $\alpha_2 = 1$. Since $\alpha_2 = 1$ in all cases $f_1(p,-1,q) = \alpha_q = \left(\frac{a+b\sqrt{-1}}{q}\right)$. Note that since $q\nmid p = \overline{B}$ we may also take $\alpha_q = \left(\frac{2(a+\sqrt{p})}{q}\right)$ and thus we recover the equality $\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4 = \left(\frac{\frac{1}{2}(a+\sqrt{p})}{q}\right)$ from above. (Alernatively we may use the relation $2(a+b\sqrt{-1})(a+\sqrt{p}) = (a+b\sqrt{-1}+\sqrt{p})^2$ from Remark 2.4, which implies that $\left(\frac{a+b\sqrt{-1}}{q}\right) = \left(\frac{2(a+\sqrt{p})}{q}\right)$ so the two statements are equivalent.)

To prove [L, Ex. 5.5, p. 176] we note that we also may write

$$\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4 = f(p,-1,q) = f_1(p,q,-1).$$

We have $A=q$, $B=p$ and $C=-1$ so $f_1(p,q,-1) = \alpha_\infty\alpha_2$. If $e^2 = pf^2+qg^2$ then $e^2-qg^2 = pf^2$ so we may take $(x,y,z) = (e,g,f)$.

We want to prove that $f_1(p,q,-1)=(-1)^{\frac{fg}{2}}\left(\frac{-1}{e}\right)$. By permuting, if necessary p and q we may assume that if $p\equiv q\equiv 5 \pmod 8$ then f is even and g is odd and if $p\not\equiv q \pmod 8$ then $p\equiv 5, q\equiv 1 \pmod 8$. We have $\overline{A}=q$, $\overline{B}=p$ so $\overline{A}\equiv\overline{B}\equiv 1 \pmod 4$ and $\overline{C}=-1$. Since $C<0$ we have $\alpha_\infty=sgn(e)$. There are two cases: $q\equiv 1 \pmod 8$ and $q\equiv p\equiv 5 \pmod 8$. In the second case by our assumption g is even. In the first case $\overline{A}\equiv 1 \pmod 8$ so $\alpha_2=\left(\frac{x+y\sqrt{\overline{A}},C}{2}\right)=\left(\frac{e+g\sqrt{q},-1}{2}\right)$. If g is even then $\alpha_2=\left(\frac{e,-1}{2}\right)\left(\frac{1+\frac{g}{e}\sqrt{q},-1}{2}\right)$. Now $\frac{1}{e}\sqrt{q}$ is an odd 2-adic integer and g is even. If $2\|g$ then $1+\frac{g}{e}\sqrt{q}\equiv 3 \pmod 4$ so $\left(\frac{1+\frac{g}{e}\sqrt{q},-1}{2}\right)=-1$. If $4\mid g$ then $1+\frac{g}{e}\sqrt{q}\equiv 3 \pmod 4$ so $\left(\frac{1+\frac{g}{e}\sqrt{q},-1}{2}\right)=1$. Hence $\left(\frac{1+\frac{g}{e}\sqrt{q},-1}{2}\right)=(-1)^{\frac{g}{2}}=(-1)^{\frac{fg}{2}}$ and so $\alpha_2=(-1)^{\frac{fg}{2}}\left(\frac{e,-1}{2}\right)$. If g is odd so f is even note that $pf^2=e^2-qf^2\equiv 1-1=0 \pmod 8$ so $4\mid f$. Now e and $g\sqrt{q}$ are odd 2-adic integers so by replacing, if necessary $\sqrt{q}$ by $-\sqrt{q}$ we have $e\equiv g\sqrt{q} \pmod 4$. It follows that $2\|e+g\sqrt{q}$. Since also $(e-g\sqrt{q})(e+g\sqrt{q})=e^2-qg^2=pf^2\vdots 16$ we get $e-g\sqrt{q}\vdots 8$ so $e+g\sqrt{q}\equiv 2e \pmod 8$. It follows that $\frac{e+g\sqrt{q}}{2e}\equiv 1 \pmod 4$ so $\left(\frac{\frac{e+g\sqrt{q}}{2e},-1}{2}\right)=1$, which implies that $\alpha_2=\left(\frac{e+g\sqrt{q},-1}{2}\right)=\left(\frac{2e,-1}{2}\right)=\left(\frac{e,-1}{2}\right)=(-1)^{\frac{fg}{2}}\left(\frac{e,-1}{2}\right)$. (Recall, $4\mid f$.) In the second case $\overline{A}\equiv\overline{B}\equiv 5 \pmod 8$ and $\overline{C}=-1$ so $\alpha_2=-\left(\frac{3x+z\sqrt{5\overline{B}},-1}{2}\right)=-\left(\frac{3e+f\sqrt{5p},-1}{2}\right)=-\left(\frac{3e,-1}{2}\right)\left(\frac{1+\frac{f}{3e}\sqrt{q},-1}{2}\right)$. Now $pf^2=e^2-qg^2\equiv 1-5\equiv 4 \pmod 8$ so $2\|f$. Since also $\frac{1}{3e}\sqrt{5p}$ is an odd 2-adic integer we have $1+\frac{f}{3e}\sqrt{5p}\equiv 3 \pmod 4$ so $\left(\frac{1+\frac{f}{3e}\sqrt{q},-1}{2}\right)=-1$. Since also $-\left(\frac{3e,-1}{2}\right)=\left(\frac{e,-1}{2}\right)$ we get $\alpha_2=-\left(\frac{e,-1}{2}\right)=(-1)^{\frac{fg}{2}}\left(\frac{e,-1}{2}\right)$. In conclusion $\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4=\alpha_\infty\alpha_2=sgn(e)\cdot(-1)^{\frac{fg}{2}}\left(\frac{e,-1}{2}\right)=(-1)^{\frac{fg}{2}}\left(\frac{-1}{e}\right)$.

Next we prove that if $p=r^2+qs^2$ then $\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4=\left(\frac{2}{q}\right)^s$ (see [L, Ex. 5.6, p. 176]). This time we use

$$\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4=f(p,-1,q)=f_2(p,q,-1).$$

We have $A=q$, $B=p$ and $C=-1$ so $ABC=-pq$. Since $p^2-pr^2=pqs^2$ we may take $(x,y,z)=(p,r,s)$. We have $f_2(p,q,-1)=\beta_\infty\beta_2$.

Since $x=p>0$ we have $\beta_\infty=1$ so we have to prove that $\beta_2=\left(\frac{2}{q}\right)^s$. Since $C=-1$ there are three cases for β_2. If $q=\overline{A}\equiv 1 \pmod 8$ then $\beta_2=\left(\frac{x,C}{2}\right)=\left(\frac{p,-1}{2}\right)=1=\left(\frac{2}{q}\right)^s$. If $q\equiv 5 \pmod 8$ and $p=\overline{B}\equiv 1 \pmod 8$ then $\left(\frac{2}{q}\right)^s=(-1)^s$ and $\beta_2=\left(\frac{2(x+y\sqrt{\overline{B}}),C}{2}\right)=\left(\frac{2(p+r\sqrt{p}),-1}{2}\right)=\left(\frac{p+r\sqrt{p},-1}{2}\right)$. If s is odd and r is even then $r^2=qs^2-p\equiv 1-5\equiv 4 \pmod 8$ so $2\|r$. We have $\beta_2=\left(\frac{p,-1}{2}\right)\left(\frac{1+r\sqrt{\frac{1}{p}},-1}{2}\right)$. But $2\|r$ and $\sqrt{\frac{1}{p}}$ is an odd integer so $1+r\sqrt{\frac{1}{p}}\equiv 3 \pmod 4$ so $\left(\frac{1+r\sqrt{\frac{1}{p}},-1}{2}\right)=-1$. It follows that $\beta_2=\left(\frac{p,-1}{2}\right)\cdot(-1)=-1=\left(\frac{2}{q}\right)^s$. (Recall that s is odd and $q\equiv 5 \pmod 8$.) If s is even and r odd then $qs^2=p-r^2\equiv 1-1=0 \pmod 8$ so $4\mid s$. Since p, $r\sqrt{p}$ are 2-adic odd integers, by multiplying $\sqrt{p}$ with ± 1 we may assume that $p+r\sqrt{p}\equiv 2 \pmod 4$, i.e. $2\|p+r\sqrt{p}$. Together with $p^2-pr^2=pqs^2\vdots 16$ we get that $8\mid p-r\sqrt{p}$ so $p+r\sqrt{p}\equiv 2p \pmod 8$ so $\frac{1}{2p}(p+r\sqrt{p})\equiv 1 \pmod 4$ so $\left(\frac{\frac{1}{2p}(p+r\sqrt{p}),-1}{2}\right)=1$, which implies that $\beta_2=\left(\frac{2p,-1}{2}\right)=1=\left(\frac{2}{s}\right)^s$ (s is even). Now assume that $p\equiv q\equiv 5 \pmod 8$, i.e. $\overline{A}\equiv\overline{B}\equiv 5 \pmod 8$, so $\beta_2=\left(\frac{-2(5x+y\sqrt{5\overline{B}}),C}{2}\right)=\left(\frac{-2(5p+r\sqrt{5p}),-1}{2}\right)=-\left(\frac{5p+r\sqrt{5p},-1}{2}\right)$. If s is odd and r even then $r^2=qs^2-p\equiv 5-5=0 \pmod 8$ so $4\mid r$ so $5p+r\sqrt{5p}\equiv 5p\equiv 1 \pmod 4$ so $\beta_2=-\left(\frac{5p+r\sqrt{5p},-1}{2}\right)=-1=\left(\frac{2}{q}\right)^s$. (Recall that s is odd and $q\equiv 5 \pmod 8$.) If s is even and r is odd then $qs^2=p-r^2\equiv 5-1=4 \pmod 8$ so $2\|s$. By multiplying $\sqrt{5p}$ with ± 1 we may assume that $2\|5p+r\sqrt{5p}$. We also have $25p^2-5pr^2=20p^2+5pqs^2\equiv 4+4\equiv 8 \pmod{16}$. ($p^2\equiv 1 \pmod 4$ so $20p^2\equiv 20\equiv 4 \pmod{16}$ and $5pq(\frac{s}{2})^2\equiv 1 \pmod 4$ so $5pqs^2\equiv 4 \pmod{16}$.) Hence $5p-r\sqrt{5p}\equiv 4 \pmod 8$ so $5p+r\sqrt{5p}\equiv 10p+4\equiv 50+4\equiv 6 \pmod 8$, which implies that $\left(\frac{5p+r\sqrt{5p},-1}{2}\right)=-1$ so $\beta_2=1=\left(\frac{2}{q}\right)^s$ (s is even).

Scholz. Assume that $p\equiv q\equiv 1 \pmod 4$ are primes such that $\left(\frac{p}{q}\right)=1$ and assume that ε is a unit of $\mathbb{Q}(\sqrt{p})$ of norm -1. Then Scholz's law states that $\left(\frac{\varepsilon}{q}\right)=\left(\frac{p}{q}\right)_4\left(\frac{q}{p}\right)_4$. (See [L, p. 167].) We could prove Scholz's law directly if $\varepsilon\in\mathbb{Z}[\left(\frac{1+\sqrt{p}}{2}\right)]\setminus\mathbb{Z}[\sqrt{p}]$. However in order to overcome the 2-adic complications we replace ε by ε^3 and we may assume that $\varepsilon = t + u\sqrt{p} \in \mathbb{Z}[\sqrt{p}]$. We have

$$\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = f(p, -1, q) = f_1(-1, p, q)$$

so $A = p$, $B = -1$, $C = q$. Since $t^2 - pu^2 = -1$ we may take $(x, y, z) = (t, u, 1)$. We have $f_1(-1, p, q) = \alpha_\infty \alpha_q \alpha_2$. But $C > 0$ so $\alpha_\infty = 1$ and $q \nmid p = \overline{A}$ so $\alpha_p = \left(\frac{x+y\sqrt{A}}{q}\right) = \left(\frac{t+u\sqrt{p}}{p}\right) = \left(\frac{\varepsilon}{p}\right)$. So we have to prove that $\alpha_2 = 1$. If $\overline{C} = q \equiv 1 \pmod 8$ then $\alpha_2 = 1$ by definition. If $\overline{C} = q \equiv 5 \pmod 8$ and $\overline{A} = p \equiv 1 \pmod 8$ then $\alpha_2 = \left(\frac{x+y\sqrt{A}, C}{2}\right) = \left(\frac{t+u\sqrt{p}, 5}{2}\right)$. But t and u have opposite parities so $t+u\sqrt{p}$ is an odd 2-adic integer, which implies that $\alpha_2 = 1$. If $p \equiv q \equiv 5 \pmod 8$, i.e. $\overline{A} \equiv \overline{C} \equiv 5 \pmod 8$ then $\alpha_2 = \left(\frac{z, 5}{2}\right) = \left(\frac{1, 5}{2}\right) = 1$.

A similar result holds if $q = 2$. Namely, if $p \equiv 1 \pmod 8$ then $\left(\frac{1+\sqrt{2}}{p}\right) = \left(\frac{2}{p}\right)_4 \left(\frac{p}{2}\right)_4$. (See [L, p. 169].) We repeat the reasoning used to prove that $\left(\frac{p}{q}\right)_4 \left(\frac{q}{p}\right)_4 = f(p, -1, q)$. By the special case 2 we have $f_2(2, p, p) = \left(\frac{2}{p}\right)_2$ and $f_2(p, 2, 2) = \left(\frac{p}{2}\right)_4$ so $\left(\frac{2}{p}\right)_4 \left(\frac{p}{2}\right)_4 = f(2, 2p, p)$. We have $f(2, 2p, p) = f(2, -2p, p) f(2, -1, p)$. But $f(2, -2p, p) = f_2(2, -2p, p)$ can be calculated using the special case 3 with $B = 2$, $C = p$. We have $n = \left(\frac{\overline{C}}{\overline{C}, 2\overline{B}}\right) = p$ and $\overline{B} = 2 \not\equiv 1 \pmod 4$ so $\beta_2 = 1$. Thus $f_2(2, -2p, p) = (-1)^{\frac{p^2-1}{8}} \cdot 1 = 1$ so $\left(\frac{2}{p}\right)_4 \left(\frac{p}{2}\right)_4 = f(2, 2p, p) = f(2, -1, p) = f_1(-1, 2, p)$. We have $A = 2$, $B = -1$, $C = p$ so $f_1(-1, 2, p) = \alpha_\infty \alpha_p \alpha_2$. We have $1^2 - 2 = -1$ so we may take $(x, y, z) = (1, 1, 1)$. Since $C = p > 0$ we have $\alpha_\infty = 1$ and since $p \nmid 2 = \overline{A}$ we may take $\alpha_p = \left(\frac{x+y\sqrt{A}}{p}\right) = \left(\frac{1+\sqrt{2}}{p}\right)$ so we have to prove that $\alpha_2 = 1$. But this follows from $\overline{C} = p \equiv 1 \pmod 8$.

References

[L] Lemmermeyer, Franz, *Reciprocity Laws: from Euler to Eisenstein*, Berlin: Springer-Verlag, 2000